

## 6 Biological Outcomes : Has customary management been successful? Discussion and Conclusions

---

### 6.1 Management Success : Discussion

The present study aimed to examine the success of customary management actions for fin-fish resources across a range of fishing pressures at different sites in Fiji and Vanuatu. In addressing the question, 'Has customary management been successful', the criteria used in this volume on biological outcomes relate to western notions of fisheries management success. Management success has not been measured against any community objectives for biological management, which were not explicitly stated.

A number of authors (Christy, 1982, Hviding and Ruddle, 1991, Pomeroy, 1994, Gadil *et al* 1993, Doulman, 1995) have proposed developing traditional management systems based on Customary Marine Tenure (CMT) combined with scientific advice. Others have questioned this notion (McCay and Acheson, 1987, Ruddle, 1998), whilst in Samoa a community management system has been established based on principles of customary tenure (King and Faasili, 1997), although that project did not examine the effectiveness of customary management. The present study addressed questions relating to the feasibility of developing customary management systems for co-management: 'Exactly how effective are existing attempts at customary management?'. Volume 2 described the effectiveness of customary management in terms of social equity and the human community. The current Volume examined customary management success in relation to the fishery resource base.

The key finding from analysis of the social outcomes of customary management was that management activities related principally to political (e.g. land tenure) and cultural requirements of the community. Management actions were not defined by the biology of the resources. Thus, management boundaries are unlikely to relate to the biological distributions of fish stocks, and management actions are unlikely to recognise key life cycle events or be based on 'scientific' criteria such as spawning seasons or size at maturity. This implies that whilst customary management interventions occur, any effects (beneficial or otherwise) on the fishery resource base are co-incidental. Resource management is not the primary aim of the customary management interventions, although it is apparent that one aim of management interventions is to amass (save-up) a 'bank' of resources for subsequent use related to social functions (see Volume 2 and 6.1.3 below). This Volume therefore investigated the status of fishery resources inside managed areas in order to establish whether customary management conferred any benefit compared to un-managed open access areas? Fishery indicators (e.g. mean length, catch rates, abundance, species assemblages) were also correlated with the level of fishing effort to see if this explained any observations.

Study sites were selected in Fiji and Vanuatu within which a number of sub-areas were identified including a managed (tabu) area and other un-managed areas. The principle customary management instruments studied at these sites were effort controls and closed areas. Gear controls have also been applied (Pellonk in Vanuatu, see Volume 2) but have generally been ignored. In Fiji explicit effort controls were applied in Verata and Naweni (both semi-commercial sites) where in order to protect native fishing rights non Vanua fishers are

excluded. In Vanuatu there are implicit effort controls in that community fishing rights areas (CFRAs) are only fished by members of the community, although some illegal fishing and poaching activities do occur by outsiders. At commercial sites in Fiji, non Vanua fishers are licensed to fish within the CFRA, but there are currently no restrictions on numbers of fishers, and thus no effort controls. Closed areas were applied in both Vanuatu and Fiji.

Management actions limit the effects of fishing (which were described in Volume 3 Chapter 1), and thus in managed areas one would expect to observe a lower level of fishing mortality (F), higher population density and biomass, and larger mean length of fish. Differences in species assemblages and species richness may also occur. It would be expected that managed areas would have low or, in the case of closed areas, zero fishing effort. It is these indicators of the status of fishery resources that were examined, inside and outside managed areas, and in relation to fishing effort in order to judge the status of the resources and draw conclusions on management success.

### **6.1.1 Are management actions respected?**

Since resource management is not the primary aim of customary management interventions, fishery management targets were not set by fishing communities in either Fiji or Vanuatu on which to judge management success. Some perception of potential biological management targets was apparently held in Vanuatu in relation to the number of fishers and extent of exploitation of the resources (see Volume 4), but this was not translated into management actions.

From the current study it was not possible to define management targets in terms of maximum catch or effort: The time-series of data gathered from the fisheries monitoring programme was insufficient to enable resource assessments; Whilst UVC studies enable the determination of mean biomass of fish for dive locations within each area, this related to a specific depth band and would not be universally applicable across the whole reef so it was not considered valid to apply such estimates to the whole reef at each area. However, in order to provide an indication of potential yield from CFRAs, it is valid to apply comparisons with similar reefs elsewhere. Dalzell (1996) indicated that typically yields from reef fisheries are in the range 4-5 t/km<sup>2</sup> per annum, but that in the Pacific yields of up to 36.9 t/km<sup>2</sup> have been reported. Tables 6.1-6.2 indicate the catch per unit area during the present study for Vanuatu and Fiji respectively, and the proportion of the potential yield this represents at 5 t/km<sup>2</sup> and 30 t/km<sup>2</sup>.

For Vanuatu, at the lower estimate of potential yield the level of catch is high, and exceeds sustainable limits in a number of areas (Table 6.1); 201, 204 in Atchin, 246T and 249 in Lelepa, 223 in Uripiv and 211 and 212 in Wala. Of the Tabu areas no catch or a very low catch was reported in Pellonk and Uripiv in 1997, and in 1998 only a very small catch was taken from 222T in Uripiv and 215T in Wala. However, at Emua and Lelepa the level of catch from tabu areas was relatively high as it was for Wala in 1997 (264T, Emua, 1998, 61.3% of lower estimate of yield; and at 246T in Lelepa, catches exceeded the lower estimate of sustainable yield in both 1997 and 1998; 215T Wala, 1997, 54.4%). As indicated in Volume 2, enforcement is highly variable. The fact that the level of catch taken from some of the areas was high is an important consideration when interpreting the results of Chapters 2-4 of this Volume. However, it does not necessarily mean in relation to the communities objectives, that tabus have not been respected (see Volume 2 and this Volume, 6.1.3). From a Western, fishery management perspective, however, we can conclude that Tabus were only respected in Pellonk, and Uripiv, and in Wala in 1998. In Wala in 1997, Emua & Lelepa,

closures were not respected. Spear gun bans in Pellonk were not respected.

The level of catch from fishing areas in Fiji was small at all of the semi-commercial sites (Table 6.2). The level of exploitation from tabu areas was comparable with that from open access areas except for 201T in Naweni in 1998 when the tabu was apparently respected and the catch was small. The level of catch from commercial sites was low to moderate, with the exception of sites 2 and 8 where relative to the lower estimate of sustainable yield, catches were high (Table 6.2). Licensing may have acted to limit entry, but no limits were applied. Note, however, that the low catches from some areas in Fiji may be due to prior overfishing.

The question remains as to whether fishing intensity would have been even greater in tabu and licensed areas in Fiji and Vanuatu had these 'management' measures not been applied.

Table 6.1. The estimated catch per annum per unit area of reef within sub-areas of study sites in Vanuatu, and the proportion of the potential yield this represents at an estimated sustainable yield of 5 t/km<sup>2</sup> and 30 t/km<sup>2</sup>. Tabu areas are indicated (T).

Site Name	Subarea	Area (sqkm)	Minimum est'd total catch pa	Minimum est'd total mt per sq km/yr	Percent est'd MSY (5 mt sq km/yr)	Percent est'd MSY (30 mt sq km/yr)
Atchin 97	200	1.351	638.69	0.47	9.46%	0.32%
Atchin 97	201	0.069	1376.08	20.06	401.25%	13.37%
Atchin 97	202	0.075	125.63	1.68	33.50%	1.12%
Atchin 97	203	0.115	186.22	1.62	32.50%	1.08%
Atchin 97	204	0.119	561.37	4.73	94.67%	3.16%
Atchin 97	205	2.090	16.00	0.01	0.15%	0.01%
Atchin 97	206	0.929		0.00	0.00%	0.00%
Lelepa 97	242	32.070	0.00	0.00	0.00%	0.00%
Lelepa 97	243	0.588	169.22	0.29	5.76%	0.19%
Lelepa 97	244	3.262	1289.77	0.40	7.91%	0.26%
Lelepa 97	245	3.472	206.20	0.06	1.19%	0.04%
Lelepa 97	246T	0.132	709.57	5.39	107.84%	3.59%
Lelepa 97	247	0.540	783.88	1.45	29.02%	0.97%
Lelepa 97	248	0.935	180.10	0.19	3.85%	0.13%
Lelepa 97	249	0.245	958.84	3.92	78.40%	2.61%
Lelepa 97	250	18.100	179.55	0.01	0.20%	0.01%
Lelepa 97	241	4.276		0.00	0.00%	0.00%
Pellonk 97	231	0.958	45.15	0.05	1.00%	0.16%
Pellonk 97	232	3.081	2244.98	0.73	14.56%	2.43%
Pellonk 97	233T		18.65	0.28	5.60%	0.93%
Uripiv 97	221	1.038	625.24	0.60	12.05%	0.40%
Uripiv 97	222T	0.107		0.00	0.00%	0.00%
Uripiv 97	223	0.194	1074.90	5.54	110.81%	3.69%
Uripiv 97	224	0.483	653.78	1.35	27.10%	0.90%
Uripiv 97	225	0.412	55.66	0.14	2.70%	0.09%
Wala 97	211	0.122	2654.28	21.74	434.77%	14.49%
Wala 97	212	0.223	1539.86	6.89	137.86%	4.60%

Site Name	Subarea	Area (sqkm)	Minimum est'd total catch pa	Minimum est'd total mt per sq km/yr	Percent est'd MSY (5 mt sq km/yr)	Percent est'd MSY (30 mt sq km/yr)
Wala 97	215 T	0.007	19.00	2.72	54.36%	1.81%
Wala 97	216	0.008	32.84	3.97	79.45%	2.65%
Wala 97	217	2.381	15.27	0.01	0.13%	0.00%
Atchin 98	200	1.351	114.32	0.08	1.69%	0.06%
Atchin 98	201	0.069	577.85	8.42	168.49%	5.62%
Atchin 98	202	0.075	111.26	1.48	29.67%	0.99%
Atchin 98	203	0.115	220.18	1.92	38.43%	1.28%
Atchin 98	204	0.119	212.29	1.79	35.80%	1.19%
Atchin 98	205	2.090	40.43	0.02	0.39%	0.01%
Atchin 98	206	0.929	233.11	0.25	5.02%	0.17%
Emua 98	262	0.667	821.45	1.23	24.65%	0.82%
Emua 98	263	1.068	1482.46	1.39	27.76%	0.93%
Emua 98	264T	0.448	1373.36	3.06	61.26%	2.04%
Lelepa 98	242	32.070	845.89	0.03	0.53%	0.02%
Lelepa 98	243	0.588	1731.62	2.95	58.95%	1.96%
Lelepa 98	244	3.262	1661.10	0.51	10.19%	0.34%
Lelepa 98	245	3.472	41.44	0.01	0.24%	0.01%
Lelepa 98	246T	0.132	1030.41	7.83	156.60%	5.22%
Lelepa 98	247	0.540	354.86	0.66	13.14%	0.44%
Lelepa 98	248	0.935	736.19	0.79	15.74%	0.52%
Lelepa 98	249	0.245	1586.72	6.49	129.74%	4.32%
Lelepa 98	250	18.100	368.98	0.00	0.00%	0.00%
Lelepa 98	241	1.065	611.96	0.11	2.15%	0.07%
Uripiv 98	221	1.038	127.76	0.12	2.46%	0.08%
Uripiv 98	222 T	0.107	29.66	0.28	5.56%	0.19%
Uripiv 98	223	0.194	731.62	3.77	75.42%	2.51%
Uripiv 98	224	0.483	140.88	0.29	5.84%	0.19%
Uripiv 98	225	0.412	124.76	0.30	6.06%	0.20%
Wala 98	211	0.122	521.25	4.27	85.38%	2.85%
Wala 98	212	0.223	2508.59	11.23	224.58%	7.49%
Wala 98	215 T	0.007		0.00	0.00%	0.00%
Wala 98	216	0.008		0.00	0.00%	0.00%
Wala 98	217	2.381		0.00	0.00%	0.00%

Table 6.2. The estimated catch per annum per unit area of reef within sub-areas of study sites in Fiji, and the proportion of the potential yield this represents at an estimated sustainable yield of 5 t/km<sup>2</sup> and 30 t/km<sup>2</sup>. Tabu areas are indicated (T).

Site Name	Subarea	Area (sqkm)	Est'd total catch pa	Est'd total mt per sq km/yr	Percent est'd MSY (5 mt sq km/yr)	Percent est'd MSY (30 mt sq km/yr)
<b>Semi-Commercial sites</b>						
Verata 96-97	16 97 T	27.95	4.31	0.15	3.08%	0.51%
Viwa (E) 96-97	17 97	29.28	1.36	0.05	0.93%	0.15%

Site Name	Subarea	Area (sqkm)	Est'd total catch pa	Est'd total mt per sq km/yr	Percent est'd MSY (5 mt sq km/yr)	Percent est'd MSY (30 mt sq km/yr)
Kubuna 96-97	18 97	85.65	26.84	0.31	6.27%	1.04%
Naweni 96-97	20 97	9.526	2.92	0.31	6.13%	1.02%
Naweni Tabu 96-97	201 97 T	8.01	3.57	0.45	8.92%	1.49%
Tacilevu 96-97	21 97	7.07	6.61	0.93	18.69%	3.12%
Verata 97-98	16 98 T	27.95	5.76	0.21	4.12%	0.69%
Viwa (E) 97-98	17 98	29.28	1.45	0.05	0.99%	0.16%
Kubuna 97-98	18 98	85.65	26.17	0.31	6.11%	1.02%
Naweni 97-98	20 98	9.526	7.50	0.79	15.74%	2.62%
Naweni Tabu 97-98	201 98 T	8.01	0.50	0.06	1.26%	0.21%
Tacilevu 97-98	21 98	7.07	5.26	0.74	14.87%	2.48%
<b>Commercial Sites</b>						
Vuda 97-98	2	55.56	358.01	6.44	128.87%	21.48%
Vitogo 97-98	5	98.89	97.43	0.99	19.70%	3.28%
Votua 97-98	6	71	18.12	0.26	5.10%	0.85%
Naviti-Marou 2 97-987		38.41	41.41	1.08	21.56%	3.59%
Waya-Naviti 97-98	8	8.81	41.55	4.72	94.33%	15.72%
Naviti-Marou 97-98	9	84.84	36.99	0.44	8.72%	1.45%
Cautata 97-98	19	3.289	3.13	0.95	19.03%	3.17%
Tavua 97-98 (inner)	121	88.23	132.74	1.50	30.09%	5.01%
Tavua 97-98 (outer)	122	98.615	230.48	2.34	46.74%	7.79%

### 6.1.2 Has management conferred any benefit compared to un-managed open access areas?

The results of UVC and univariate and multivariate analyses of fisheries monitoring data for Vanuatu and Fiji are summarised in Tables 6.3-6.4 respectively which compare resource indicators from managed and open access areas.

- Neither fisheries nor UVC data indicated any differences in mean length of fish in closed or licensed areas compared to open access for Fiji or Vanuatu.
- Mortality estimates were inadequate for comparison
- Species assemblages were not significantly different in closed and open areas of Fiji and Vanuatu from UVC or fisheries data from MDS analysis except for gill net caught fish in 1998 in Vanuatu. Fishery data indicated that the species assemblage was most different for 222 (Uripiv) and 264 (Emua) compared to open access areas, which is consistent with the observation that the level of fishing and catch from 246 (Lelepa) was high. The finding for Emua is perhaps surprising given that Table 6.1 indicated that considerable fishing had also occurred inside that tabu area. As in the example of Lelepa, this apparent anomaly may be explained by the pattern of resource use in the Emua closed area (see Volume 2, and below).
- Species assemblages at commercial Fiji sites were weakly correlated to number and cost of licences.
- Univariate UVC and fisheries analyses indicated some benefits to management with increased abundance (and cpue) of certain families / guilds or overall in Wala, Uripiv (UVC) and Lelepa (Fisheries) in Vanuatu, and in Naweni in 1998 in Fiji (UVC and

fisheries). The finding for increased catch rate (an indicator of abundance) in the tabu area in Lelepa (246) was surprising given the high catches taken from the tabu area (see Table 6.1). However, this is apparently explained by the pattern of resource use (see Volume 2 and 6.1.3 below).

- Such increases in abundance were not related to growth of fish (no size differences), and thus may indicate an aggregation effect in less disturbed areas. The behaviour of fish subjected to consistent fishing may also change, affecting their catchability.

Table 6.3. Summary results of analyses of indicators of the status of fishery resources in open access and managed (*tabu*) areas in Vanuatu, and correlation of these indicators with fishing effort and abundance.

<b>Analysis performed</b>	<b>Overall Observations</b>	<b>Tabu areas vs open access areas</b>
<b>UVC analyses</b>		
Habitat	Habitat differences occurred between areas and sites from the MDS ordination, but ANOSIM indicated these were not significant.	MDS showed that two tabu areas (246T at Lelepa and 263T at Emua) differed from open access areas, but ANOSIM showed the differences to be insignificant. Habitat was comparable in studied closed and open areas.
Species assemblages	<p>Certain species, family and trophic group differences in abundance occurred between sites and between areas within sites. At family and trophic group level there was a weak trend for increasing biomass at low levels of effort, but this was not shown to be significant across areas except for Serranidae and Kyphosidae with ln transformed data.</p> <p>Across dive-sites a significant negative trend for biomass with effort occurred for Balistidae, Lutjanidae, Serranidae and Siganiidae only. Multiple regression indicated that habitat variables played a more significant role in determining fish abundance than fishing effort.</p>	<p>Univariate analyses indicated that differences in species abundance were not consistent inside and outside tabu areas: some species were more abundant in tabu areas, but the reverse was also true. At family and trophic level, Lethrinids were more abundant in 215 T (Wala), as were planktivores; Lutjanidae and Mullidae were more abundant in 222 T at Uripiv as were piscivores. These differences in biomass were consistent with expectation for the level of fishing at each area. No differences between closed and open areas occurred in Lelepa or Emua at family or trophic level.</p> <p>Multivariate analyses with MDS suggested that 263T in Emua differed from the open access sites in Emua, but ANOSIM indicated no significant differences for any closed areas vs open areas. Open area 247 in Lelepa differed from other areas, but the Tabu area 246T did not differ.</p>
Mean length		Wala, Uripiv, Lelepa, Emua - no differences in mean length between closed and open areas for any species.

<b>Analysis performed</b>	<b>Overall Observations</b>	<b>Tabu areas vs open access areas</b>
<b>Univariate analyses</b>		
Mean catch rate by area (index of abundance)	Within sites few significant differences occur in cpue. The differences were greater across sites	Data were available for 3 tabu areas 222 (Uripiv), 246 (Lelepa) and 264 (Emua). Only 246 had a significantly higher catch rate than open access areas within the relevant site.
Correlation of catch rate with fishing intensity	Catch rate data were poorly correlated with fishing intensity	Relative to the level of fishing effort, area 222 had low/expected catch rates, those at 264 were as expected, those at 246 tended to be high.
Mean length of key species by area	Few differences in mean length between sub areas occurred for fish caught within sites with any gear. there were significant differences between sites.	No significant differences in mean length occurred for fish caught in tabu areas compared to open access areas.
Correlation of mean length with fishing intensity and abundance	Although not ubiquitous, a significant negative correlation existed between mean length and fishing intensity, and positive correlation with abundance for some species and gears.	Mean length of fish in tabu areas varied, but did not differ from expectation.
Correlation of fishing mortality of key species with fishing intensity and abundance	Total mortality was not correlated to fishing intensity or abundance except for <i>Lethrinus harak</i>	Insufficient data were available except for <i>Ctenochaetus striatus</i> at 246, where total mortality was low.
<b>Multivariate analyses</b>		
Species assemblages	No correlation existed between species assemblages and fishing intensity, reef area or distance from landing site. These variables were strongly inter-correlated.	No significant differences in species assemblage occurred between tabu and open access areas, except for fish caught with gill nets in 1997/8. Species assemblages were most different at 222 and 264, whilst at 246 they were more similar to other areas.

Table 6.4. Summary results of analyses of indicators of the status of fishery resources in open access and managed (*tabu*) areas in Fiji, and correlation of these indicators with fishing effort and abundance.

Analysis performed	Site status	Overall Observations	Tabu areas vs open access areas
<b>UVC analyses</b>			
Habitat		Habitat differences between sites were not significant, but significant differences occurred between areas across sites : Tacilevu differed from all other areas, but those within Tavua and Naweni were similar	Tabu areas did not show significant differences from open access areas.
Species assemblages		Certain species, family and trophic group differences in abundance occurred between sites and between areas within sites, but fishing effort did not contribute significantly to explaining variation in mean biomass. In Tavua inshore reefs (121) had greater biomass than offshore reefs. MDS plots indicated offshore reefs separated from inshore reefs at this site.	Uni-variate tests indicated significant differences in invertebrate feeding rates between open access areas and tabu areas, but not conclusive.  Multivariate tests showed significant differences between open access area 20 and tabu areas, but differences were not significant.
Mean length		Two parrot fishes, <i>Clorurus bleekeri</i> and <i>C. sordidus</i> , had greater mean lengths on inshore reefs in Tavua compared to offshore reefs.	No significant differences in mean length were detected in the closer to shore areas.
<b>Uni-variate analyses</b>			
Mean catch rate by area (abundance)	Semi-commercial		No difference between 201 (Naweni) in 1997/8 and 1998/9
	Commercial		No differences in catch rates between 201 (Naweni) in 1997/8 and 1998/9
Correlation of catch rate with fishing intensity and abundance	All sites	No correlation with any gear/year combination	
	Semi-commercial	No correlation except for gillnets in 1997/8	No consistent trend for catch rates when 16, 201 were compared to other sites
	Commercial	No correlation with any gear/year combination	No consistent trend for catch rates when 16, 201 were compared to other sites
Mean length of key species by area	Semi-commercial	Few differences in mean length by area were observed, but fish caught at 18 and 21 tended to be larger.	No evidence that length of fish caught in those in open access areas was significantly larger than those in tabu areas
	Commercial sites	No consistent differences in mean length within or across commercial sites, except fish at 122 (offshore reef) which were larger	No consistent trend for length of fish caught in those in open access areas
	Semi-commercial vs commercial sites	Fish caught by handlines from commercial sites were larger than those from semi-commercial sites.	No consistent trend for length of fish caught in those in open access areas

<b>Analysis performed</b>	<b>Site status</b>	<b>Overall Observations</b>	<b>Tabu areas vs open</b>
Correlation of mean length with fishing intensity and abundance	All sites	Few species indicated significant correlations between mean length and fishing intensity and abundance, and except in two cases observations were inconsistent with expectation. Environmental differences or fishing practices (especially between commercial / semi-commercial areas) may have a stronger influence than fishing intensity.	Mean length of fish in
Correlation of fishing mortality of key species with fishing intensity and abundance		For some species, fishing mortality was positively correlated to fishing intensity and negatively correlated to abundance	Limited data was available for species from tabu areas consistent with the length for one observation on
<b>Multivariate analyses</b>			
Species assemblages	All sites	No correlation existed between species assemblages and fishing intensity or reef area.	
	Semi-commercial sites	Species assemblage was not correlated to fishing intensity. There was no apparent difference in species assemblage across all areas, but this could not be statistically tested.	Species assemblage significantly different
	Commercial sites	Species assemblage was not correlated to fishing intensity.	Species assemblage licences per area, an
	Semi-commercial vs commercial sites	Family assemblages caught by handlines were significantly different between commercial and Semi-commercial sites.	

### 6.1.3 Explaining the observations

Tables 6.3-6.4 compare resource indicators from managed and open access areas and correlate them with fishing effort in order to explain observations reported in Section 6.1.2 above. The key findings were:

- Mean length, mortality, abundance and species assemblages at Fiji and Vanuatu showed no, or only weak correlation with fishing effort. The abundance of few species, families or guilds was correlated to effort.
- Managed areas were generally consistent with expectation relative to the level of effort applied in them, with few exceptions.
- Such differences as occurred could be attributed to other factors such as habitat (Vanuatu), inshore or offshore reefs (Fiji), or level of commercialisation (Fiji)

Mean length was not significantly correlated with fishing intensity or abundance, and such variations as did occur were believed to relate to environmental or fishing practice differences between areas and fishing sites. This was especially the case in Fiji where there was a division between semi-commercial and commercial fishing sites and fish from the latter were larger. Fishing mortality was not correlated with fishing intensity or abundance in Vanuatu, and weakly correlated for certain species in Fiji. The mortality of fish from Tabu areas was consistent with expectation however.

Species assemblages were examined in both UVC and multivariate studies of fishery data. Species assemblage was not correlated to fishing intensity in either Fiji or Vanuatu for either UVC or fishery data. Jennings and Polunin (1996) in a similar UVC study in Fiji found that at very low levels of fishing effort, species assemblages were significantly different, but that at higher levels of effort there was no difference. The fact that species assemblage and fishing effort were not correlated in the present study suggests that fishing effort at all the study sites in Vanuatu and Fiji was above a threshold level at which these changes in species assemblage occur. Tabu areas were subject to significant fishing also. However, it will be recalled that gill net data for 1998 did indicate species assemblage differences between tabu and open access areas in Vanuatu. The species assemblage in Lelepa (246) where high catches were taken, did not differ markedly from open access areas, whilst that in Uripiv (222, low catch) and Emua (264, high catch) did.

Catch rate data as an indicator of abundance indicated little correlation with fishing effort in either Vanuatu or Fiji. Catch rates in tabu areas were generally consistent with expectation for the level of fishing applied. Exceptions were 246T at Lelepa in Vanuatu, where catch rates were higher than expected for the level of effort, and 201T in 1998 at Naweni in Fiji.

Thus, few indicators of the status of resources revealed differences between managed and open access areas (Section 6.1.2). Benefits of closed areas were apparent where effort was low, and resulted in higher abundance of fish recorded at the family and trophic level, when making direct comparisons between tabu and open access areas. However, generally the value of indicator variables for managed areas were consistent with expectation for the level of effort applied in them. This suggests that where closures may be enforced, and effort controls applied, management benefits will occur. Furthermore, however, there remained cases of where management had apparently conferred some additional benefit:

- Species assemblages at commercial Fiji sites were weakly correlated to number and cost of licences. This assumes that the resulting species assemblage is beneficial -

In fact, this relates to targeting effects by commercial fishers rather than effort limitation (species assemblage was not correlated with effort), and the benefits to the community would be reduced competition for reef fish. Volume 2, chapter 2 indicates the markedly different species composition taken by commercial and semi-commercial fishers in Fiji;

- Species assemblages in 222 in Emua (Vanuatu) differed from open access areas despite high catches from that area (this assumes that the resulting species assemblage is beneficial - in terms of conservation and bio-diversity this is the case, but no analysis was performed of the relative value of the different catches likely to result from these different species assemblages);
- Catch rates in 246 in Lelepa (Vanuatu) and 201T in Naweni in 1998 (Fiji) exceeded expectation relative to the level of effort.

In relation to the closed areas, the observations may be explained by the pattern of resource use (see Volume 2). In Vanuatu, the community utilise the resources in the 'closed' area in a different way from those in open access areas, which may be fished at any time. In the closed areas, apart from one illegal invasion in area 246 in Lelepa, the tabus were respected by the communities of Emua and Lelepa, with the intention of allowing the resource to 'build up'. Then at special events (such as the visit by the Prime Minister who was from Lelepa, or Christmas, for example) the community will fish the resource. Thus pulsed fishing occurs and apparently higher catch rates were achieved at Lelepa. This relates to the fact that any aggregation or increased catchability effects result in high catch rates initially. Since only a short pulse of fishing occurs, catch rate decreases are not apparent and catch rates are not averaged over a long time period as they are for the open access areas. Hence relative to the level of effort, the catch rate is high. The effect of the pulsed fishing at Emua was a different species assemblage. In relation to this site, it should be noted that surrounding areas had also been closed prior to the study, and the aggregate effect of these closures may have resulted in the benefits described. In the case of Naweni in Fiji, the catches recorded occurred after re-opening of the closure. This indicates that short term closures lead to short term measurable benefits above expectation for the level of effort. Longer term benefits would not be expected.

## **6.2 Management success : Conclusions**

Customary management measures that were studied related to short term closures and effort controls (unrestricted licensing).

### ***Closed areas***

Polunin (1984) referring to Indonesia and Papua New Guinea, suggested that conservation is not the intent of many traditional closed areas and that the resulting management actions are the outcome of conflict over scarce resources. Similarly in Fiji and Vanuatu, conservation was not the primary aim, and management actions were related to land tenure, cultural and political issues rather than to western notions of sustainable resource use. The results of this study have indicated that limited evidence for the benefits of customary management can be shown from the western perspective. The status of fishery resources inside tabu areas was generally consistent with the level of exploitation of those areas. Benefits from fishery closures were not comparable to those reported by other authors in relation to long term marine protected areas (e.g. Watson et al 1998, Russ and Alcala, 1998. See also Chapter 1, this Volume). However, benefits from short term closure were detected at two levels:

- Closures resulted in reduced effort in tabu areas, and it was shown that abundance and catch rates were higher at low levels of fishing effort.
- The pattern of use of closed areas resulted in short term benefits to the community through increased catch rates and species assemblages.

Apart from the fact that closures studied were short term, the limited evidence for benefits of management also relate to the size and location of tabu areas (no striking differences were observed in terms of abundance, size of fish, mortality of fish or species assemblages and richness inside tabu areas compared to open access areas). Frequently these were very small (see Volume 2) and were located to maximise political and cultural aims rather than in relation to the distribution of resources.

From these findings, it would appear that whilst customary management, due to its short term nature, does not result in the long term benefits associated with western or centrally managed long term marine protected areas, it does result in short term benefits. These benefits satisfy the community objectives of 'saving up' resources for social functions. Providing that the period that fishing is permitted within such areas is limited, it may be that such a pattern of resource use could provide long term benefits for overall resource sustainability, and provide short term benefits to the community. The short term benefits also act to provide an incentive to maintain the closure during the remainder of the year. Further investigation is warranted to establish if such forms of closure are an appropriate model for closed area management.

### ***Effort controls and Licensing***

Effort controls were exercised at three levels:

- Protection of native fishing rights ie. exclusion of non community members from fishing within Customary fishing rights areas (explicit, as in semi-commercial sites in Fiji, or implicit as in Vanuatu, see Volume 2 and 6.1.1);
- Licensing of non community members to fish within CFRAs, as in commercial sites in Fiji, although there are currently no restrictions on the number of fishers;
- Closed areas, which, in the way they have been applied in Fiji and Vanuatu, restrict effort for part of the year.

Gear controls would also act to limit effort but in the study site where these were nominally applied (Pellonk, Vanuatu), they were not enforced.

The results of this study have indicated that limited evidence for the benefits of customary management and effort controls can be shown from the western perspective. Explanatory variables were only weakly correlated with fishing effort and the more obvious differences between areas could be attributed to other factors such as habitat (Vanuatu), inshore or offshore reefs (Fiji), or level of commercialisation (Fiji). Nevertheless, managed areas were generally consistent with expectation relative to the level of effort applied in them. Species assemblages at commercial Fiji sites were weakly correlated to number and cost of licences.

In relation to effort controls, despite the weak correlation, the fact that the results of this study were consistent with expectation in relation to the level of fishing implies that where effort limitation is practical and can be enforced, then it can be a useful management tool.

The lack of evidence for substantive fishery and conservation benefits from customary management actions, when measured against Western values and objectives for management, is consistent with a similar study of customary management in Kenya by Mclanahan *et al* (1997). That study indicated that 'Presently, traditional management (*in Kenya*) is not effective in protecting species diversity or ecological functions, which was probably never the intention of the customs'. Some of the most degraded reefs in Kenya are those subject to customary management, whilst centrally administered national parks have performed significantly better. That study also highlights the dangers of researchers with different academic foci (e.g. biologists and anthropologists who have different views on human organisation and its effect on common resources) working independently, and stresses that a multidisciplinary approach is required as was the case with the present study.

### **Concluding Remarks**

Little evidence for the success of customary management against western objectives of the long term benefits to resource sustainability could be demonstrated from this study. Short term benefits, which meet community objectives were, however, apparent. Both Fiji and Vanuatu are signatories to the United Nations Convention on the Law of the Sea and the FAO code of Conduct for Responsible Fisheries which, from a national perspective, define the obligations of governments related to conservation, management and development of natural resources (see Mees, 1999). Thus governments need to ensure that fishery resources are managed for long term sustainability, not just short term gain. The customary management practices (or, forms of 'human-resource organisation') applied at study sites in Fiji and Vanuatu are compatible with national policies and could be adapted to achieve long term benefits. The community institutional structure for resource management exists. Thus given this situation, there is a role for government to provide scientific advice in a co-management partnership in order to enable sustainable resource management. Government and NGOs would be involved particularly in:

- Ensuring an integrated approach so that marine resource management does not occur in isolation from that of other activities in the coastal zone and beyond;
- Co-ordinating co-management activities involving all the stakeholders, including interactions with resource users outside the CFRAs;
- Identifying sites most at risk from overfishing and resource degradation, in order to target these for immediate management action - Government management actions should be pro-active rather than reactive, and should be based on the best available information and not deferred until more is available;
- Working closely with fishing communities through the extension services and providing a training curriculum for those extension workers;
- Generating public awareness and education programmes to inform the public and resource stakeholders on issues relating to resource sustainability;
- Providing scientific advice to resource custodians on appropriate management actions, noting that for small scale artisanal fisheries in particular, some areas should be closed to fishing to provide refuges for fish stocks and to protect habitats. In the context of the study locations this could include, for example, advice on factors such as the optimal size, location, number and duration of closures to achieve conservation goals; on optimum gear sizes and deployment; and on licensing strategies to optimise revenue and limit effort; on the scale of management area appropriate for different resources;
- Undertaking targeted research and pilot scale adaptive management / demonstration

projects in order to enable the provision of appropriate management advice, and test its application;

- Implementing a monitoring programme, and one suitable at the community level;
- Ensuring legal support for areas under tenure and for any management actions including enforcement by communities.

Communities would be involved in:

- Defining, in the light of advice, management actions for the different resources or CFRAs - management actions need not be the same within each CFRA (particularly for sedentary species), but for certain resources, such as migratory pelagic species, it may be appropriate for common management actions, and for co-ordinated, joint management action amongst CFRAs, requiring co-operation amongst resource custodians;
- Providing the institutional framework for management;
- Implementing and, within appropriate limits, enforcing management actions;
- Working co-operatively with adjacent fishing communities to jointly develop management strategies
- Monitoring the fishery and effects of management.

There are a number of implications to developing a system of co-management, not least the implicit and potential change in emphasis of management actions to promote their biological function above that of their socio-political functions. These are discussed more fully in Volume 2 and in the Guidelines for co-management, Volume 5. Whilst there are clearly limitations on the extent to which communities will change in order to adopt human-resource centred organisational patterns rather than, or in parallel with human-social (self) organisational patterns, cultural systems in Fiji and Vanuatu are dynamic, and have in the past adapted to changing circumstances. This suggests that there is good potential to develop resource management strategies, mutually acceptable to fishing communities and national government, based on the principle of co-management in relation to areas of customary marine tenure.



## 7 References Cited

---

- Adams, T.J.H. (1993) The Regulation of Fiji's fisheries 1987-1992. SPC, Noumea, New Caledonia.
- Adams, T.J.H. and P.J.Dalzell (1994) Artisanal Fishing. Paper presented at the East-West Center Workshop on Marine Biodiversity Issues in the Pacific Islands. Univ. of Hawaii.
- Adams, T.J.H. (1996) Governance of fisheries and aquaculture in the Pacific Islands region. Review paper for the 3<sup>rd</sup> Dialogue on the ACP-EU Research Initiative, Belize, December, 1996.
- Allison, G.W., J. Lubchenco and M.H. Carr (1998) Marine reserves are necessary but not sufficient for marine conservation. *Ecological Applications*, 8(1), Supplement 1998, pp. 579-592.
- Amos, M. (1993) Traditionally based marine management systems in Vanuatu. *Traditional Resource Management and Knowledge Information Bulletin* No. 2, K. Ruddle (ed.). The South Pacific Commission, Noumea, New Caledonia, 1993, pp.14-17.
- Ayling, A.M. and A.L. Ayling (1986). A biological survey of selected reefs in the Capricorne Section of the Great Barrier Reef Marine Park. *Rep. Great Barrier Reef Mar. Park Authority*, pp.1-61. GBRMPA, Townsville, Australia.
- Ayling, A.M., A.L. Ayling and B.D. Mapstone (1992). Possible effects from fishing pressure on recruitment rates of the coral trout (*Plectropomus leopardus*: Serranidae) *In* Hancock, D.A (ed) Australian Society for Fish Biology - Workshop - Recruitment Processes, Hobart, 21 Aug. 1991. Australian Government Publishing Service :16, pp. 210-215.
- Beddington, J.R. (1984) The response of multispecies systems to perturbations. *In*: May RM (ed) *Exploitation of Marine Communities*. Springer, Berlin, pp. 209-225.
- Beddington, J.R., and J.G. Cooke (1982) Harvesting from a prey-predator complex. *Ecological Modelling*, 14, pp.155-177.
- Beddington, J. R. and J. G. Cooke (1983). The potential yield of fish stocks. *FAO Fisheries Technical Paper*. 242: 47pp.
- Bohnsack, J.A. (1982) Effects of piscivorous predator removal on coral reef fish community structure. *In* Caillet, G.M and C.A. Simstead (eds), *Gutshop '81: Fish food habits studies*. Wash. Sea Grant Publ., Seattle, pp. 258-267.
- Bohnsack, J.A. (1998) Application of marine reserves to reef fisheries management. *Australian Journal of Ecology*, 23 No 3, pp. 293-304.

- Bonnemaison, J. (1992) Social and Cultural Aspects of Land Tenure. In: Lamour, P. Land Tenure in Vanuatu.
- Bray, J. R. and J.T. Curtis (1957). An ordination of the upland forest communities of Southern Wisconsin. *Ecol. Monogr.* 27, pp.325-349.
- Bromley, D.W.(1992) Introduction. In *Making the commons work - Theory, Practice and Policy*. ed. D.W. Bromley, ICS Press, San Francisco, 1992. 319pp.
- Carr, M.R. (1997) *PRIMER User Manual*. Plymouth UK: Plymouth Marine Laboratory.
- Carrier, J.G (1987) 'Marine Tenure and Conservation in Papua New Guinea: Problems of Interpretation', in B.J. McCay and J.M. Acheson (eds) *The Question of the Commons*, pp.142-167, Tuscon: University of Arizona Press.
- Christy, Jr., F.T. (1982) Territorial use rights in marine fisheries : Definitions and conditions. *FAO Fish Tech Pap* (227).
- Clarke, K. R. (1993). Non-parametric multivariate analyses of changes in community structure. *Aust. J. Ecol.* 18, pp.117-143.
- Clarke, K. R. and Warwick, R. M. (1994). Change in Marine Communities. An Approach to statistical analysis. Natural Environment Research Council, UK, 144 pp.
- Cohen, F.G. (1989). Treaty Indian tribes and Washington State: The evolution of tribal involvement in fisheries management in the U.S. Pacific Northwest. In : Pinkerton, E. (ed.) *Co-operative management of local fisheries. New directions for improved management and community development*. UBC Press, Vancouver.
- Cooke, A. (1994) *Qoliqoli* of Fiji - Some preliminary research findings in relation to management. In *Traditional marine tenure and sustainable management of marine resources in Asia and the Pacific* : Proceedings of the International Workshop held at the University of the South Pacific, Suva, Fiji, 4-8 July, 1994. G.Robin South, D. Goulet, S. Tuqiri & M. Church (eds). International Ocean Institute, Marine Studies Programme, University of the South Pacific, Fiji, 1994.
- Cooke, J.G. and J.R. Beddington (1984) The relationship between catch rates and abundance in fisheries. *IMA Journal of Mathematics Applied in Medicine and Biology* 1, pp.391-405.
- Cordell, J. and M.A. McKean (1982) Sea Tenure in Bahia, Brazil. In: Bromley, D.W. (ed.) (1992) *Making the commons work: theory, practice and policy*. Institute for Contemporary Studies Press, San Francisco, California.
- Craik, W.J.S. (1981a) Recreational fishing on the great barrier reef. *Proc. Int. Coral Reef Symp., 4th* 1, pp.47-52.
- Craik, W.J.S. (1981b) Underwater survey of coral trout *Plectropomus leopardus* (Serranidae) populations in the Capricornia section of the Great Barrier Marine Park. *Proc. Int. Coral Reef Symp., 4th* 1, pp.53-58.

- Dale, N. (1989). Getting to co-management: Social learning in the redesign of fisheries management. In : Pinkerton, E. (Ed.) Co-operative management of local fisheries. New directions for improved management and community development. UBC Press, Vancouver.
- Dalzell, P. (1992) Deep Reef Slope Fishery Resources of the South Pacific : a summary and analysis of the dropline fishing survey data generated by the activities of the SPC Fisheries Programme between 1974 and 1988. SPC Tech Doc. Inshore Fisheries Research Project, No 2.
- Dalzell, P. (1996) Reef fisheries yields *In* N.V.C. Polunin and C.M. Roberts (eds) Tropical Reef Fisheries, Chapman and Hall.
- David, G. and E. Cillaurren (1992). National fisheries development policy for coastal waters, small-scale fishing, and food self-reliance in Vanuatu. *Man and Culture in Oceania* 8: 35-38.
- Doherty, P.J. and P.F. Sale (1985) Predation on juvenile coral reef fishes: An exclusion experiment. *Coral Reefs* 4, pp.225-234.
- Doherty, P.J. and D. McB. Williams (1988) The replenishment of coral reef fish populations *Oceanogr. Mar. Biol.* 26, pp.487-551.
- Doulman, D.J. 1993. Community-based fishery management. Towards the restoration of traditional practices in the South Pacific. *Marine Policy*, Vol.17, No. 2.
- Fairburn, T.I.J. (1992). Traditional reef and lagoon tenure in Western Samoa and its implications for giant clam mariculture. P.169-189 In: Tisdell, C. (ed.). *Giant Clams in the Sustainable Development of the South Pacific*. Canberra, ACIAR.
- FAO 1995. Code of conduct for responsible fisheries. Rome, FAO, 41pp.
- FAO 1995. Precautionary approach to fisheries : Part 1. Guidelines on the precautionary approach to capture fisheries and species introductions. Elabor

ated by the Technical Consultation on the Precautionary Approach to Capture Fisheries (including species introductions). Lyskil, Sweden, 6-13 June 1995. FAO Fisheries

- Filipsson, O. (1989). The effects of fisheries on the size of fishes in mountain lakes. *Inf. Soetvattenlab.*, Drottningholm. No. 3, pp1-18.
- Fong, G. (1994), Case study of a traditional marine management system : Sasa village, Macuata Province, Fiji. *FAO Field Report RAS/92/TO5*. No. 94/1, Rome, FAO, 1994.
- Gadil, M, F. Berkes and C. Folke (1993) Indigenous knowledge for biodiversity conservation. *Ambio*, 22, No. 2-3, pp.151-156.
- Gayanillo, F.C., P.Sparre and D.Pauly (1994) FiSAT The FAO-ICLARM Stock Assessment Tools (FiSAT) User's Guide. *FAO Computerised Information Series, 7, FAO Rome / ICLARM Contribution No. 1048*.
- Guenette, S., T. Lauck and C. Clark (1998) Marine reserves: From Beverton and Holt to the present. *Reviews in Fish Biology* 8, pp.251-272.
- Gulland, J.A. (ed) (1977) Fish population dynamics: the implications for management. Wiley, 422p.
- Gulland, J.A. (ed) (1979) Report of the FAO/IOD workshop on the fishery resources of the western Indian Ocean south of the equator. *IOFC/Dev./79/45*.
- Gulland, J.A. (1983) Fish stock assessment. A manual of basic methods. FAO/Wiley Series in Food and Agriculture. Vol 1. Wiley, Chichester.
- Hassett, D. (1994). The community-based approach to natural resource management - A Review prepared for the Traditional Management of Artisanal Fisheries Project, CEMARE, Univ. of Portsmouth.
- Holmes, P. (1996) Land Tenure in Vanuatu. Custom, Culture, Tradition.... and Development? MSc Thesis, Australian National University, Canberra. (Unpubl).
- Hviding, E. (1988). Marine tenure and resource development in Marovo Lagoon, Solomon

Islands: traditional knowledge, use and management of marine resources, with implications for contemporary development. FFA Report No. 88/35, Forum Fisheries Agency, Honiara.

- Hviding, E. and K.Ruddle (1991) A regional assessment of the potential role of customary marine tenure (CMT) systems in contemporary fisheries management in the South Pacific. FFA Report 91/71, South Pacific Forum Fisheries Agency, Honiara, Solomon Islands.
- Hviding, E. and G.B.K. Baines (1994) Community-based fisheries management, Tradition and the challenges of development in Marovo, Solomon Islands. *Development and Change* Vol.25 (1994), pp13-39. Institute of Social Studies.
- Jennings, S., E.M. Grandcourt, N.V.C. Polunin (1995) The effects of fishing on the diversity, biomass and trophic structure of Seychelles' reef fish communities. *Coral Reefs* 14, pp.225-235.
- Jennings, S. and J.M. Lock (1996) Population and ecosystem effects of fishing. In N.V.C. Polunin and C.M. Roberts (eds) *Tropical Reef Fisheries*, Chapman and Hall, pp. 193-218.
- Jennings, S. and M.J. Kaiser (1998) The effects of fishing on marine ecosystems. *Advances in Marine Biology*, 34, pp.202-352.
- Jennings, S., S.S. Marshall and N.V. C. Polunin (1995) Seychelles Marine Protected Areas : Comparative structure and status of reef fish communities. *Biol. Conservation* 75, pp. 201-209.
- Jennings, S. and N.V.C. Polunin (1995) Biased underwater visual census biomass estimates for target species in tropical reef fisheries. *Journal of Fish Biology* 47, pp. 733-736.
- Jennings, S. and N.V.C. Polunin (1996) Effects of fishing effort and catch rate upon the structure and biomass of Fijian reef fish communities. *Journal of Applied Ecology* 33, pp. 400-412.
- Jentoft, S. and B. McCay (1995) User participation in fisheries management: Lessons drawn from international experience. *Marine Policy*, Vol.19, No. 3, pp. 227-246.
- Johannes, R.E. (1981). *Words of the Lagoon. Fishing and Marine lore in the Palau District of Micronesia*. Berkeley: University of California Press.
- Johannes, R.E. (1994) Government-supported, village-based management of marine resources in Vanuatu. *FFA Report 94/2*, South Pacific Forum Fisheries Agency, Honiara, Solomon Islands, 1994.
- King, M and U. Faasili (1997) Community based management of fisheries and the marine environment. Pacific science Association Inter-congress, July 1997, 9 pp.
- Klien, M. (1992). Effect of exploitation on growth, age-structure, and recruitment of an underfished population of European whitefish. *Biology and Management of Coregonid Fishes 1990*. Todd T.N. & M. Luczynski [eds.] Vol. 39, no.3-4, pp.807-815.

- Koslow, J.A., F. Hanley, and R. Wicklund (1988) Effects of fishing on reef fish communities at Pedro Bank and Port Royal Cays, Jamaica. *Mar. Ecol. Prog. Ser.* 43, pp.201-212.
- Kulbicki, M., G. Mou Tham, P.Thollot and L. Wantiez (1993) Length-weight relationships of fish from the lagoon of New Caledonia. *Naga* 2&3, pp. 26-30.
- Kulbicki, M. and L. Wantiez (1990) Variations in the fish catch composition in the Bay of st Vincent, New Caledonia, as determined by experimental trawling. *Aust. J. Mar. Freshwat. Res.* 41(1), pp.121-144.
- Kunatuba, P. (1989). Inshore fisheries development and management: the South Pacific experience. In: Report on the Workshop on Customary Tenure, Traditional Resource Management and Nature Conservation, South Pacific Regional Environment Programme, Noumea, March 1988.
- Lamour, P., (ed.) (1984) Land Tenure in Vanuatu. University of the South Pacific, Institute of Pacific Studies.
- Lindley, R. (199 ) End of Contract Report of the Fisheries Advisor to the Fisheries Department.
- Lock, J.M. (1986) Fish yields of the Port moresby barrier and fringing reefs. *Tech. Rep. Dep. Primary Ind. (Papua New Guinea)* 86(2), pp.1-17.
- McCay, B. and J.M. Acheson (1987) Human ecology of the commons, Chapter 1 pp 1-34 In: *Making the commons work - Theory, Practice and Policy.* ed. D.W. Bromley, ICS Press, San Francisco, 1992. 319pp.
- McCay, B.J. (1988). Muddling through the clam beds: Cooperative management of New Jersey's hard clam spawner sanctuaries. *Journal of shellfish Research.* Vol. 7, No.2, pp.327-340.
- McClanahan, T.R. and N.A. Muthiga (1988) Changes in Kenyan coral reef community structure and function due to exploitation. *Hydrobiologia* 166, pp.269-276.
- McClanahan, T.R., H. Glaesel, J. Rubens and R. Kiambo (1997) the effects of traditional fisheries management on fisheries yields and the coral reef ecosystems of southern Kenya. *Environmental conservation*, 24 (2), pp.105-120.
- Mees C.C. (1999) The monitoring control and surveillance of domestic artisanal fisheries. In: Report of a regional workshop on fisheries monitoring control and surveillance, Kuala Lumpur and Kuala Terengganu, Malaysia, 29 June - 3 July 1998, Supplement 2, Technical Papers, PP93-118. FAO/Norway Government Co-operative Programme - GCP/INT/648/NOR, FAO Rome, February 1999.
- MRAG (1993) A review of coral reef fisheries stock assessment literature - Fish Management Science Programme, Overseas Development Administration, London 43pp.
- MRAG (1996) Management of Multispecies Tropical Fisheries - project final technical report to ODA for the Fisheries Management Science Programme, Project R5484, MRAG London, 193 pp.

- Munro, J.L. (Ed. 1983) Caribbean Coral Reef Fishery Resources. A second edition of 'The biology, ecology, exploitation and management of Caribbean coral reef fishes : scientific report of the ODA/UWI fisheries ecology research project, 1969 - 1973: University of the West Indies, Jamaica'. *ICLARM Studies and Reviews 7*, 276p. ICLARM, Manila.
- Oakerson, R.J. (1992) Analyzing the commons: A Framework. In *Making the commons work - Theory, Practice and Policy*. ed. D.W. Bromley, ICS Press, San Francisco, 1992. 319pp.
- Ostrom, E. (1994) Institutional analysis, design principles and threats to sustainable community governance and management of commons. In *Community management and common property of coastal fisheries in Asia and the Pacific: concepts, methods and experiences*. ICLARM Conf. Proc. 45, Manila, Philippines, 1994.
- Pauly, D. (1979) Theory and management of tropical multispecies stocks. A review with emphasis on the Southeast Asian Demersal fisheries. *ICLARM Studies and Reviews No. 1*, 35pp ICLARM Manila.
- Pauly, D. (1980). On the interrelationships between natural mortality, growth parameters and mean environmental temperature in 175 fish stocks. *Journal du Conseil, Conseil International pour l'Exploration de la Mer*. 39(3), pp.175-192.
- Pauly, D. (1987a). A review of the ELEFAN system for analysis of length-frequency data in fish and aquatic invertebrates, p. 7-34. In D. Pauly and G.R. Morgan [eds.] *Length-based methods in fisheries research*. ICLARM Conference Proceedings 13, 468 p. ICLARM, Manila, Philippines.
- Pido, M.D., R.S. Pomeroy, M.B. Carlos and L.R. Garces (1996) A handbook for rapid appraisal of fisheries management systems (version 1). *ICLARM Education Service 16*, ICLARM, Makati, Philippines, 1996.
- Pinkerton, E. (1989). *Co-operative management of local fisheries. New directions for improved management and community development*. UBC Press, Vancouver.
- Polunin, N.V.C. (1984) Do traditional marine reserves conserve? A view of Indonesian and New Guinean evidence. In: *Maritime institutions in the western Pacific*, ed. K. Ruddle and T. Akimichi, pp. 267-283 Osaka, Japan, National Museum of Ethnology.
- Polunin, N.V.C (1997) Technical report on the UVC component of the Project, based on field work in Fiji and Vanuatu in 1996, and analysis in Newcastle.
- Polunin, N.V.C. and S.Jennings (1998) Differential effects of small-scale fishing on predatory and prey fishes on Fijian reefs. pp 95-124 In: *Dynamics of Tropical Communities* (ed D.M. Newbery, H.H.T. Prins, N Brown), Symposia of the British Ecological Society 33. London: Blackwell
- Polunin, N.V.C. and C.M. Roberts (1993) Greater biomass and value of target coral-reef fishes in two small Caribbean marine reserves. *Marine Ecology – Progress Series* 100, pp.167-176.

- Pomeroy, R.S. (ed.) 1994. Community management and common property of coastal fisheries in Asia and the Pacific: concepts, methods and experiences. ICLARM Conference Proceedings. 45, Manila, Philippines.
- Raja, N.M., Viswanathan, K.K. and R.S.Pomeroy (1998) Fisheries Co-management and Transaction Costs. NAGA, July-September, 1998. ICLARM. Philippines.
- Randall, J.E., G.R. Allen and R.C. Steene (1990) Fishes of the Great Barrier Reef and Coral Sea. Bathurst, NSW, Australia: Crawford Press.
- Roberts, CM (1998) Sources sinks and the design of marine reserve networks. *Fisheries* 23(7), pp.16-19.
- Roberts, C.M. and N.V.C. Polunin (1991) Are marine reserves effective in management of reef fisheries? *Reviews in Fish Biology and Fisheries*, 1, pp. 65-91.
- Ross, M.R. and G.A. Nelson (1992). The influence of stock abundance and water temperature upon the growth dynamics of haddock and yellowtail flounder on Georges Bank. Copenhagen, Denmark, *ICES 1992*. 15pp.
- Ruddle, K., E. Hviding and R.E. Johannes (1992) Marine resources management in the context of customary tenure. *Marine Resource Economics* 7(4), pp. 275-296.
- Ruddle, K.E. (1998) The context of policy design for existing community based fisheries management systems in the Pacific Islands. *Ocean and Coastal Management* 40 (2-3), pp.105-126.
- Russ, G.R. (1984) Effects of fishing and protective management on coral reefs at four locations in the Visayas, Philippines (phase II). *UNEP-NRMC Coral reef Monitor. Project, Natural Resource Management Centre, Philippines*, pp.1-54.
- Russ, G.R. (1985) effects of protective management on coral reef fishes in the central Philippines. *Proc. Int. Coral Reef Congr.* 5th 4, pp.219-224.
- Russ, G.R. (1989). Distribution and abundance of coral reef fishes in the Sumilon Island reserve, central Philippines, after nine years of protection from fishing. *Asian Mar. Biol.* 6, pp.59-71.
- Russ, G.R. (1991) Coral reef fisheries : effects and yields. In : *The Ecology of Fishes on Coral Reefs* Ed. P. F. Sale, Academic Press, San Diego.
- Russ, G.R. and A.C. Alcala (1989) effects of intense fishing pressure on an assemblage of coral reef fishes. *Mar ecol. Prog. Ser.* 56, pp.13-27.
- Russ, G.R. and A.C. Alcala (1998a) Natural fishing experiments in marine reserves 1983-1993: roles of life history and fishing intensity in family responses *Coral Reefs*, 17(4), pp. 399-416
- Russ, G.R. and A.C. Alcala (1998b) Natural fishing experiments in marine reserves 1983-1993: Community and trophic responses, 17(4), pp. 383-397.

- Samoilys, M.A. (1988) Abundance and species richness of coral reef fish on the Kenyan coast: the effects of protective management and fishing. *Proc. Int. Coral Reef Symp. 6th* 2, pp.261-266.
- Samoilys, M. (1992) Review of the underwater visual census method developed by the QDPI/ACIAR project: Visual assessment of reef fish stocks. Conference and Workshop Series QC92006, Department of Primary Industries, Brisbane.
- Samoilys, M. and G.M. Carlos (1992) *Development of an underwater visual census method for assessing shallow-water reef fish stocks in the south-west Pacific*. Queensland Department of Primary Industry, Cairns, Australia, 1992.
- Sen, S. and J.R. Nielsen (1996) Fisheries co-management: a comparative analysis. *Marine Policy*, Vol.20, No. 5, pp. 405-418.
- Schaap, A. and R. Green (1988). Fish Communities on reefs subjected to different levels of fishing pressure. *Tech. Rep. Mar. Lab. Dep. Sea. Fish. Tasm.* No. 31, 44pp
- Shulman, M.J., J.C. Ogden, J.P. Ebersole, W.N. McFarland, S.L. Miller and N.G. Wolf (1983) Priority effects in the recruitment of juvenile coral reef fishes. *Ecology* 64: pp.1508-1513.
- Sokal, R.R. and F.J. Rohlf (1995). *Biometry: the principles and practice of statistics in biological research* (3rd ed.). W.H. Freeman and Company, New York. 887p.
- Sphigel, M, and L. Fishelson (1991) Experimental removal of piscivorous groupers of the genus *Cephalopholis* (Serranidae) from coral habitats in the Gulf of Aqaba (Red Sea). *Env Biol Fish* 31: pp.131-138.
- Thompson, R. and J.L. Munro (1983a). The biology, ecology and bionomics of Caribbean reef fishes: Lutjanidae (snappers). pp. 94-109. In *J.L. Munro (ed.) Caribbean coral reef fishery resources. ICLARM Stud. Rev* 7.
- Townsley, P.G. (1993) *A manual on rapid appraisal methods for coastal communities*. Bay of Bengal Programme, Madras, India, 1993.
- Wantiez, L., P. Thollot and M. Kulbicki (1997) Effects of marine reserves on coral reef fish communities from five islands in New Caledonia. *Coral Reefs* 16(4) pp.215-224.
- Watson, M.D. and R.F.G. Ormond (1994) Effect of an artisanal fishery on the fish and urchin populations of a Kenyan coral reef. *Mar. Ecol. Prog. Ser.* 109: pp.115-129.
- Watson, M.D., T. Righton, and R.F.G Ormond (1998) The effects of fishing on coral reef fish abundance and diversity. *J. Mar. Biol. Ass. UK* 76, pp.229-233.

Personal Communications

Volume 2

Vanuatu -

Belden Ham, Atchin Island community, North Malekula  
Ben Norman, Emua Village, Efate Island  
David Kalorip, Wala Island community, North Malekula  
Dorestay Kenneth, Fisheries Department, Port Vila  
Douglas Meto, Lelepa Island community, West Efate  
Francis Hickey, Nasonal Kuljural Senta blong Vanuatu, Port Vila  
Gary Preston, Gillett, Preston and Associates, Noumea, New Caledonia  
George Plant, National Planning Office, Port Vila  
Kalmasai Kalsakau, SMET, Port Vila, Efate  
Leah Horowitz, Australia National University, Canberra, ACT  
Moses Amos, Director of Fisheries, Port Vila  
Ralph Regenvanu, Nasonal Kuljural Senta blong Vanuatu, Port Vila  
Silas Nicholson, Atchin Island community, North Malekula

Fiji -

Aimutti Sadiq, Kavoli Landing, Tavua *Qoliqoli*  
Alena Tinai, Naweni Village, Vanua Levu  
Apenisa Botilagi, Tacilevu Village, Vanua Levu  
Felix Poni, Fiji Fisheries Division, Lautoka, Viti Levu  
Joe Radrodro, Fiji Fisheries Division, Lautoka, Viti Levu  
Joeli Veitayaki, MSP, USP, Suva  
Krishna Swarmy, SFO R & D, Fisheries Division, Lami, Viti Levu  
Maria Draulele, Tacilevu Village, Vanua Levu  
Native Fisheries Commission, Suva, Viti Levu  
Ratu Ovini Bokini, Tavualevu Village, Viti Levu  
Ratu Tevita Temaipeau, Fisheries Division, Savusavu, Vanua Levu  
Ratu Tulala, Ucunivanua Village, Viti Levu  
Robin South, Director, MSP, USP, Suva  
Rusiate Botilagi, Tacilevu Village, Vanua Levu  
Sairusi (Fisheries Liaison Officer, Tavua *Qoliqoli*, Tavualevu Village, Viti Levu  
Semisoni, SFO Central Division, Fiji Fisheries Division, Nausori, Viti Levu  
Siri Wakatibau, Naweni Village, Vanua Levu  
Suresh Chand, SFO, Fiji Fisheries Division, Lautoka, Viti Levu  
Timoci Tavusa, Fiji Fisheries Division, Lami, Viti Levu  
Tony Chamberlain, MSP, USP, Suva  
Vena Ram Bidesi, MSP, USP, Suva  
Vekila Vuki, MSP, USP, Suva